

Short Note

Anthracen-9-ylmethylene-(3,4-dimethylisoxazol-5-yl)amine

Abdullah M. Asiri ^{1,2,*} and Salman A. Khan ¹

¹ Chemistry Department, Faculty of Science, King Abdulaziz University, P.O. Box 80203, Jeddah, Saudi Arabia

² The Center of Excellence for Advanced Materials Research, King Abdulaziz University, Jeddah, P.O. Box 80203, Saudi Arabia

* Author to whom correspondence should be addressed; E-Mail: aasiri2@kau.edu.sa; Fax: +966 2 6952292.

Received: 8 September 2011 / Accepted: 13 September 2011 / Published: 13 September 2011

Abstract: The title compound, anthracen-9-ylmethylene-(3,4-dimethylisoxazol-5-yl)amine (**3**), was synthesized in high yield by reaction of anthracene-9-carbaldehyde and 5-amino-3,4-dimethylisoxazole in ethanol. The structure of this new compound was confirmed by elemental analysis, IR, ¹H NMR, ¹³C NMR and GC-MS spectral analysis.

Keywords: Schiff base; anthracene aldehyde; isoxazole

Five-membered heterocyclic compounds with special reference to isoxazole derivatives are an important subset of the natural products that exhibit biological activities, including anticancer [1], antifungal and antibacterial properties [2]. These compounds have also found application in electroluminescent devices and UV stabilization polymers [3]. Isoxazole-containing Schiff bases can form complexes with d4, d5, d6 metals, such as zinc, cadmium, cobalt, nickel, osmium and iridium, they exhibit luminescent properties and have been widely investigated in recent experiments [4]. They are applicable in the fields of materials science such as non-linear optics (NLO) [5], optical limiting [6], electrochemical sensing and Langmuir films [7]. In the present paper, we are reporting a novel isoxazole-containing Schiff base from the reaction of anthracene-9-carbaldehyde with 5-amino-3,4-dimethylisoxazole [8].